

Finite-temperature properties of the two-orbital Anderson model

Luis Craco

Instituto de Física Gleb Wataghin - UNICAMP, C.P. 6165, 13083-970 Campinas - SP, Brazil
(February 1, 2008)

The metallic phase of the two-orbital Anderson lattice is study in the limit of infinite spatial dimensions, where a second order perturbation treatment is used to solve the single-site problem. Using this approximation, in the Kondo regime, we find that the finite temperature properties of the conduction electrons exhibit the same behaviour as observed in the metallic phase of the two-channel Kondo lattice. Possible connections between these two models are discussed.

71.27.+a, 75.30Mb, 71.10.Fd

The normal metallic state of a number of three-dimensional heavy-fermion compounds can be quite well described by the Landau Fermi liquid theory. At zero temperature (T) the heavy-fermion quasi-particles have an infinite lifetime and a large effective mass. As a consequence of this *coherent* regime the heavy-fermion compounds present at low T a very large electronic specific heat coefficient $\gamma(T) = C(T)/T$.¹ The physical properties of these materials are related to strongly correlated electrons in $4f/5f$ orbitals, and a proper model Hamiltonian for the description of these properties is the Anderson Hamiltonian.¹

The Fermi liquid regime in the single impurity Anderson model can be identified by the behaviour of the f -electron self-energy $\Sigma^f(\omega)$ at low temperatures near of the Fermi level (ε_F). The f -electron self-energy in the single impurity Anderson model is $\mathbf{k}$ -independent and respects the Fermi liquid requirements.² Performing perturbation expansion in the f -electron Coulomb interaction (U) at low T and near ε_F Yamada showed that $\text{Re}\Sigma^f(\omega) \approx -\omega$ and $\text{Im}\Sigma^f(\omega) \approx -(\omega^2 + (\pi T)^2)$.³ For the lattice case and in the infinite dimensional limit, where $\Sigma^f(\omega)$ is also $\mathbf{k}$ -independent,^{4,5} it was shown by Georges *et al.*⁶ that the metallic phase of the Periodic (one-orbital) Anderson model can be a Fermi liquid. The behaviour of the real and imaginary part of $\Sigma^f(\omega)$ in $d = \infty$ was studied by Schweitzer and Czycholl⁷ by a self-consistent second order perturbation in U . In this case the imaginary part of $\Sigma^f(\omega)$ vanishes near ε_F in accordance with the Luttinger theorem, and the real part has a negative slope in the same region.

The investigations presented above were done only for one-channel versions of the Anderson Hamiltonian. The multichannel one impurity Anderson model, where a localised f -electron hybridises with several conduction bands (orbitals), was studied in the large U limit.^{8,9} In this limit the model shows non-Fermi liquid behaviour. A non-Fermi liquid behaviour is also present in the multichannel Kondo impurity problem. This problem was introduced by Nozières and Blandin,¹⁰ and the exact solution was obtained by Andrei and Destri¹¹ and Wiegmann and Tsvelick¹² in terms of the Bethe *Ansatz*. The non-Fermi liquid behaviour in the multichannel Kondo impurity model comes from the overcompensation of the Kondo spins by the conduction electrons.¹³ The over-

compensation mechanism was also used to explain the non-Fermi liquid behaviour in the two-channel Kondo (tCK) lattice^{14,15} and the two-channel Anderson (tCA) lattice¹⁶ in the Kondo limit.

The conclusion about the non-Fermi liquid regime in the tCK lattice was made by Jarrell *et al.*¹⁴ looking at the one-particle properties of the conduction electrons. In this case, the real part of the self-energy of the conduction electrons $\Sigma^c(\omega)$ present a positive slope near the Fermi energy. The imaginary part of $\Sigma^c(\omega)$ instead going to zero in a quadratic way as in Fermi liquid systems it goes away from zero as $\omega, T \rightarrow 0$. Despite of these properties the system is metallic and the single-particle density of states (DOS) of the conduction electrons has a finite value at the Fermi level. This metallic regime (*incoherent metal*) was used to explain the physical properties of a number of heavy Fermion compounds where the Fermi liquid paradigm can not be applied.^{14,18} For example, the incoherent metal regime have been used to explain the unusual resistivity of UBe_{13} .¹⁶

In addition, in Ref. [14] the tCK model was studied by means of quantum Monte Carlo (QMC) simulation in the limit of high dimensions. It is well known that QMC method provides very accurate results at intermediate and high temperatures. However, it does not provide any information about the explicit form of the self-energy of the problem. For example, in Jarrell's *et al.*¹⁴ work the single particle self-energy was obtained by inverting the relation $G^c(\omega) = D(\omega - \Sigma^c(\omega))$ ¹⁷ for different temperatures. It must be noted that the nonanalytic form of $\Sigma^c(\omega)$ was postulated from these numerical results. Thus, in order to understand the origin of the incoherent metal regime in tCK lattice, it is important to find out a model which describes these novel properties and allow us to obtain an explicit form for $\Sigma^c(\omega)$. Here, we address precisely this problem: the study of a multi-orbital model which correctly takes into account the incoherent behaviour of the tCK lattice and provides an explicit form for the self-energy of the conduction electrons of each orbital.

In this letter we study a multi-orbital Anderson lattice (*mOA*) model in the high dimension ($d = \infty$) limit. We discuss the formal exact solution of this multi-orbital Hamiltonian as well as the finite temperature properties of the two-orbital version of this problem. In this case a

second order perturbation treatment is applied to solve the impurity problem in the presence of U . As we have mention before, we find that the conduction electrons show the same behaviour as those obtained by Jarrell *et al.*¹⁴ for the tCK lattice.

The m OA model consists of the usual f -electron Hamiltonian of the periodic Anderson lattice model, m identical orbitals of noninteracting c -electrons and the local hybridisation between f and c electrons. The complete Hamiltonian can be written as

$$H = -\frac{t^*}{2\sqrt{d}} \sum_{\langle ij \rangle \sigma \alpha} c_{i\alpha\sigma}^\dagger c_{j\alpha\sigma} + E \sum_{i\sigma} n_{i\sigma}^f + U \sum_i n_{i\uparrow}^f n_{i\downarrow}^f + \sum_{i\sigma\alpha} V_{\alpha\sigma} \left(c_{i\alpha\sigma}^\dagger f_{i\sigma} + f_{i\sigma}^\dagger c_{i\alpha\sigma} \right), \quad (1)$$

where $c_{i\alpha\sigma}^\dagger$ ($c_{i\alpha\sigma}$) creates (destroys) a conduction electron on site i and orbital $\alpha = 1, 2, \dots, m$ of spin σ , and $f_{i\sigma}^\dagger$ ($f_{i\sigma}$), creates (destroys) a localised f -electron on site i of spin σ . The sites i form an infinite dimensional hypercubic lattice and the hopping is limited to the nearest neighbours. The scaled hopping integral $t^* = 1$ determines the energy unit. The hybridisation term $V_{\alpha\sigma}$ is site independent, but it can have different values for different orbitals or spin directions.

To obtain the formal exact solution of the m OA model it is convenient to define a new set of conduction electron operators: $\{a_{i1\sigma}^\dagger, a_{i2\sigma}^\dagger \dots a_{im\sigma}^\dagger\}$. In this case $a_{i1\sigma}^\dagger \equiv 1/\bar{V}_\sigma \sum_\alpha V_{\alpha\sigma} c_{i\alpha\sigma}^\dagger$ and the remainder $a_{i\alpha\sigma}^\dagger$ operators are written in order to preserve the definition of $a_{i1\sigma}^\dagger$ and the fermion commutation relations. The normalisation factor is $\bar{V}_\sigma = \sqrt{\sum_\alpha V_{\alpha\sigma}^2}$.

In the new representation the m OA Hamiltonian (Eq. (1)) is written as

$$H = -\frac{t^*}{2\sqrt{d}} \sum_{\langle ij \rangle \sigma \alpha} a_{i\alpha\sigma}^\dagger a_{j\alpha\sigma} + E \sum_{i\sigma} n_{i\sigma}^f + U \sum_i n_{i\uparrow}^f n_{i\downarrow}^f + \sum_{i\sigma} \bar{V}_\sigma \left(a_{i1\sigma}^\dagger f_{i\sigma} + f_{i\sigma}^\dagger a_{i1\sigma} \right). \quad (2)$$

It is clear from Eq. (2) that the multi-orbital problem for the f -electrons is reduced to a one-orbital problem with a renormalised hybridisation $\bar{V}_\sigma$. As the local approximation for the Anderson (one-orbital) lattice is exact in the limit of infinite dimensions,^{5,19} the formal exact solution for the f -electron one-particle Green's function in $d \rightarrow \infty$ is given by⁵

$$G_{\mathbf{k}\sigma}^f(i\omega_n)^{-1} = i\omega_n - \Sigma_\sigma^f(i\omega_n) - \frac{\bar{V}_\sigma^2}{i\omega_n - \epsilon_{\mathbf{k}}}. \quad (3)$$

As we intend in this work to clarify the physical origin of the incoherent properties of the tCK model, the conduction electron Green's function of each particular orbital must be known. In the context of the mapped Hamiltonian (Eq. (2)) the α -orbital Green's function

$G_{ij\sigma}^{\alpha c}(\tau) \equiv -\langle \hat{T} c_{i\alpha\sigma}(\tau) c_{j\alpha\sigma}^\dagger(0) \rangle$ is obtained when the $c_{i\alpha\sigma}$ operators are written in terms of the $a_{i\alpha\sigma}$ operators, and it is straightforward to show that

$$G_{ii\sigma}^{\alpha c}(i\omega_n) = \left(1 - \frac{V_{\alpha\sigma}^2}{\bar{V}_\sigma^2} \right) D(i\omega_n) + \frac{V_{\alpha\sigma}^2}{\bar{V}_\sigma^2} D(i\omega_n - \bar{V}_\sigma^2 \mathcal{G}_\sigma), \quad (4)$$

where $\mathcal{G}_\sigma(i\omega_n)^{-1} \equiv i\omega_n - \Sigma_\sigma^f(i\omega_n)$ and $D(z)$ is the Hilbert transform of the uncorrelated density of states of the conduction band $\rho_0(\epsilon)$.¹⁷ For a hypercubic lattice in $d = \infty$ $\rho_0(\epsilon) = (1/\sqrt{\pi}) \exp(-\epsilon^2)$.

The two terms of Eq. (4) can be easily understood from the transformation discussed above. The first term of the right-hand side of this equation describes the contributions of the $(m-1)$ free orbitals and the second term is related to the one-orbital Anderson problem of Eq. (2). It is important to notice that the formal exact solution of the m OA model (Eqs. (3) and (4)) is completely general concerning the number of orbitals as well as the values of the hybridisation on different orbitals.

In principle, it is possible to assume that the *scattering processes* in each orbital can be described by means of a self-energy.^{14,19} By definition, the α -orbital Green's function can be written in terms of the self-energy of the conduction electrons $\Sigma_{\mathbf{k}\sigma}^{\alpha c}(i\omega_n)$ as

$$G_{\mathbf{k}\sigma}^{\alpha c}(i\omega_n)^{-1} \equiv i\omega_n - \epsilon_{\mathbf{k}} - \Sigma_{\mathbf{k}\sigma}^{\alpha c}(i\omega_n). \quad (5)$$

In order to recover Eq. (4) by summing Eq. (5) over momentum, the self-energy of the α -orbital conduction electrons must be given by

$$\Sigma_{\mathbf{k}\sigma}^{\alpha c}(i\omega_n) = V_{\alpha\sigma}^2 \mathcal{G}_\sigma(i\omega_n) \times \left\{ 1 + \frac{(\bar{V}_\sigma^2 - V_{\alpha\sigma}^2) \mathcal{G}_\sigma(i\omega_n)}{i\omega_n - (\bar{V}_\sigma^2 - V_{\alpha\sigma}^2) \mathcal{G}_\sigma(i\omega_n) - \epsilon_{\mathbf{k}}} \right\}. \quad (6)$$

The requirement that in the $d \rightarrow \infty$ limit the local interactions give rise to a $\mathbf{k}$ -independent self-energy for the f -electrons is satisfied for the m OA lattice (see Eq. (3)). However, a $\mathbf{k}$ -dependent self-energy for the α -orbital conduction electrons is obtained.

The first term of $\Sigma_{\mathbf{k}\sigma}^{\alpha c}(i\omega_n)$, $V_{\alpha\sigma}^2 \mathcal{G}_\sigma(i\omega_n)$, is the self-energy of one-orbital Anderson lattice. This term is completely local and describes the scattering processes in the α -orbital. The second term is relevant only for multi-orbital systems, where $\bar{V}_\sigma^2 \neq V_{\alpha\sigma}^2$. As $(\bar{V}_\sigma^2 - V_{\alpha\sigma}^2)$ is related to all other orbitals different from α , the $\mathbf{k}$ -dependent term can be considered as an *effective correction* to the one-orbital self-energy. This effective correction describes the scattering process that happens in the remaining orbitals.

Different methods can be used to solve the single site problem in the $d = \infty$ limit.^{5,20} One of these methods is the iterative perturbation theory (IPT).²¹ IPT shows to be a good approximation for describing the Fermi liquid properties of the Hubbard model.²² For the particular case of the periodic Anderson model the IPT results

are in good agreement with exact diagonalisation⁵ and QMC.²³ The IPT method is normally applied for values of $U/t^* < 3$, however it has the property of correctly taking into account the limit of $V \rightarrow 0$.⁵ This allows us to study the one-particle properties of the m OA lattice near the Kondo limit.




FIG. 1. One-particle properties of the c -electrons for $m = 2$, $V = 0.3535$, $U = 2.0$ and three different temperatures. (a) Density of states, (b) Real and (c) Imaginary parts of the self-energy, respectively.

Let us now turn our attention on the spectral properties of the conduction electrons of the two-orbital Anderson (tOA) model. The one-particle properties of f -electrons will not be considered here because they are well known from studies of the periodic Anderson model.^{5,7,23} In our study we have chosen $U = 2.0$ and $V = 0.3535$. Note that $V = 0.3535$ for $m = 2$ means $\bar{V} = 0.5$.

In Fig. 1-a we display the single particle density of states (DOS) for the conduction electrons of tOA model. The DOS has a finite value at the Fermi level for all temperatures. Such metallic behaviour is related to charge fluctuations and to the contribution of the non-hybridised electrons. The $\mathbf{k}$ -independent real and imaginary part of the c -electrons self-energy $\Sigma^c(\omega) \equiv V^2 \mathcal{G}(\omega)$ (see Eq. (6)) is plotted in Fig. 1-b and 1-c, respectively. As one can see in Fig. 1-b, the real part of the self-energy exhibits a positive slope near the Fermi level and this slope decreases with increasing temperature. It is well known that for a Fermi liquid system the slope of $Re\Sigma^c(\omega)$ must be negative. Therefore, as it was pointed out in Ref. [14] the positive slope observed in Fig. 1-b describes the breakdown of the quasiparticle concept. Non-Fermi liquid properties can be also observed in the imaginary part of the self-energy (see Fig. 1-c). From this figure it is clear that $Im\Sigma^c(\omega)$ does not approach to the Fermi liquid form $Im\Sigma(\omega) \approx -(T^2 + \omega^2)$ as $\omega, T \rightarrow 0$. Note that similar behaviour as presented in Fig. 1 has been observed in

the one-particle properties of the conduction electrons of tCK lattice.¹⁴

It is worth noticing that the incoherent properties shown in Fig. 1-b and 1-c are closely related to the definition of $\Sigma^c(\omega)$. Concerning this point, we follow the idea of Jarrell *et al.*¹⁴ where the self-energy is defined in order to account for the Hilbert transform of the Green's function of the conduction electrons in each channel. It is well known that $\Sigma^f(\omega)$ (see Eq. (3)) shows Fermi liquid behaviour at low temperatures near to the Fermi level.⁷ From Eq. (6) one can easily conclude that the Fermi liquid properties of the f -electrons imply incoherent properties for the conduction electrons of the periodic Anderson model as well as the m OA model. Hence, to decide whether a model of two or more different particles is in a coherent regime or not, it is important to consider the behaviour of all particles.




FIG. 2. DOS of c -electrons at $T = 0.20$ for $m = 2$ (solid line) and $m = 1$ (dot-dashed line).

In Fig. 2 we compare the DOS for the conduction electrons of the tOA model with those of the periodic Anderson model. In this figure we consider two different values of hybridisation: $V = 0.3535$ ($m = 2$) and $V = 0.5$ ($m = 1$), such that the correlation effects are taken into account at the same level for both systems. This allows us to study the effect of the non-hybridised particles in the two-orbital system. For $T = 0.20$ the contribution of this particles completely suppresses the hybridisation gap and a flat region in the DOS of $m = 2$ is observed near the Fermi level. Despite the differences in the Hamiltonian parameters and the method used to solve the impurity problem similar flat behaviour have been reported in Ref. [14] for tCK model.

Let us now proceed to understand the similarities between our results and those of tCK model.¹⁴ In the Kondo limit the m OA model (Eq. (1)) is mapped onto a multi-orbital Kondo (m OK) model. In order to show this result one can apply the resolvent perturbation theory²⁴ in the subspace defined by the $a_{i1\sigma}$ and f operators, see Eq. (2). If the hybridisation is the same for both spin directions

the mOK Hamiltonian is written as

$$H_{Kondo} = -\frac{t^*}{2\sqrt{d}} \sum_{\langle ij \rangle \sigma \alpha} c_{i\alpha\sigma}^\dagger c_{j\alpha\sigma} + k \sum_{i\alpha} V_\alpha^2 \mathbf{S}_i \cdot \mathbf{s}_{i\alpha} \\ + k \sum_{i\alpha \neq \alpha'} V_\alpha V_{\alpha'} [S_i^z (c_{i\alpha\uparrow}^\dagger c_{i\alpha'\uparrow} - c_{i\alpha\downarrow}^\dagger c_{i\alpha'\downarrow}) \\ + S_i^+ c_{i\alpha\downarrow}^\dagger c_{i\alpha'\uparrow} + S_i^- c_{i\alpha\uparrow}^\dagger c_{i\alpha'\downarrow}], \quad (7)$$

where $k \equiv U/(|E|(|E| + U))^1$ and $\mathbf{S}_i, \mathbf{s}_{i\alpha}$ are the Kondo and the α -orbital conduction electron spin operators, respectively. Note that Eq. (7) is precisely the Schrieffer-Wolf transformation of Eq. (1).

The first two terms on the right hand side of Eq. (7) can be considered as the simplest generalisation of the *one* and *two-channel* Kondo models into a multi-channel one. The remaining terms are related to non-diagonal exchange processes between the impurity and the conduction electrons in the different orbitals. The relevant contributions of these non-diagonal processes will appear only in fourth order in the hybridisation term, see Eq. (6). It is clear that the fourth order contributions act only as small corrections in the Kondo limit, $(kV_\alpha V_{\alpha'})^2 \ll 1$. This explain the agreement between our results and those of tCK model.

Summarising, the simplest extension of the periodic Anderson model into a multi-orbital Anderson (mOA) model is introduced for the first time in this letter. The model is studied in the limit of high dimensions, where a second order perturbation treatment (IPT) is used to solve the impurity problem. Using this approximation at finite temperatures we find that the single-particle properties of the conduction electrons in the Kondo regime for the two-orbital Anderson model are the same as those for the two-channel Kondo lattice. We have explained this agreement in terms of the irrelevance of the non-diagonal exchange processes in the Kondo limit. Finally, we wish to point out the importance of our results since they open a new possibility of study the mCK problem in the limit of high dimensions.

ACKNOWLEDGMENTS

The author wishes to acknowledge, respectively, M. A. Gusmão and R. Bulla for useful discussions in the early stages of this work. It is also a pleasure to acknowledge M. Foglio for useful comments. This work has been partially supported by the Brazilian agency Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Max-Planck-Institut für Physik Komplexer Systeme (MPIPKS), and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). Part of IPT study was performed in the CRAY Y-MP2E of the Centro Nacional de Supercomputação da Universidade Federal do Rio Grande do Sul, at Porto Alegre, Brazil.

- ¹ P. Fulde, in *Electron Correlations in Molecules and Solids*, edited by P. Fulde in Springer Series in Solid-State Sciences Vol. 100 (Springer-Verlag, 1991).
- ² D. L. Cox and N. Grewe, Z. Phys. B **71**, 321 (1988); A. C. Hewson Phys. Rev. Lett. **70**, 4007 (1993).
- ³ K. Yamada Prog. Theor. Phys. **53**, 970 (1975).
- ⁴ W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989); see also E. Müller-Hartmann, Z. Phys. B **74**, 507 (1989).
- ⁵ A. Georges, G. Kotliar, W. Krauth and M. Rozenberg, Rev. Mod Phys. **68**, 13 (1996).
- ⁶ A. Georges, G. Kotliar and Q. Si, Int. J. Mod. Phys. B **6**, 705 (1992).
- ⁷ H. Schweitzer and G. Czycholl, Z. Phys. B **79**, 377 (1990).
- ⁸ D. L. Cox and A. E. Ruckenstein, Phys. Rev. Lett. **71**, 1613 (1993).
- ⁹ J. Kroha, P. Wölfe and T. A. Costi, Phys. Rev. Lett. **79**, 261 (1997).
- ¹⁰ P. Nozières and A. Blandin, J. Phys. (Paris) **41**, 193 (1980).
- ¹¹ N. Andrei and C. Destri, Phys. Rev. Lett. **52**, 364 (1984).
- ¹² A. M. Tsvelick and P. B. Wiegmann, Z. Phys. B **54**, 201 (1984).
- ¹³ P. Schlottmann and K. Lee, Phys. Rev. B **52**, 6489 (1995).
- ¹⁴ M. Jarrell, H. Pang, D. L. Cox and K. H. Luk, Phys. Rev. Lett. **77**, 1612 (1996).
- ¹⁵ M. Jarrell, H. Pang and D. L. Cox, Phys. Rev. Lett. **78**, 1996 (1997).
- ¹⁶ F. B. Anders, M. Jarrell and D. L. Cox, Phys. Rev. Lett. **78**, 2000 (1997).
- ¹⁷ $D(z) \equiv \int_{-\infty}^{+\infty} d\epsilon \rho_0(\epsilon)/(z - \epsilon)$ is the Hilbert transform of the uncorrelated density of states of the conduction band.
- ¹⁸ D. L. Cox, Physica B **223-224**, 453 (1996).
- ¹⁹ R. Consiglio and M. A. Gusmão, Phys. Rev. B **55**, 6825 (1997).
- ²⁰ L. Craco and M. A. Gusmão, Phys. Rev. B **54**, 1629 (1996).
- ²¹ M. J. Rozenberg, G. Kotliar and X. Y. Zhang, Phys. Rev. B **49**, 10181 (1994); H. Kajueter and G. Kotliar, Phys. Rev. Lett. **77**, 131 (1996).
- ²² X. Y. Zhang, M. J. Rozenberg and G. Kotliar, Phys. Rev. Lett. **70**, 1666 (1993).
- ²³ M. Jarrell *et al.*, Phys. Rev. Lett. **70**, 1670 (1993).
- ²⁴ H. Keiter and G. Morandi, Physics Reports **109**, 227 (1984).
- ²⁵ J. R. Schrieffer and P. A. Wolff, Phys. Rev. **149**, 491 (1966).